

The Japanese 55-year Reanalysis JRA-55

--- progress and status ---

Y.Harada, S.Kobayashi, Y.Ota, M.Moriya, S.Yasui, K.Onogi, A.Ebita, H.Kamahori, C.Kobayashi, H.Endo, K.Miyaoka, R.Kumabe, and K.Takahashi

Japan Meteorological Agency (JMA)



Japanese Reanalysis

1st JRA-25

By JMA and CRIEPI



CRIEPI:
 Central Research Institute of Electric Power Industry

2nd JRA-55 By JMA

JRA-55 Nickname

→ JRA Go! Go!







JRA-25 (ni-go)



Contents



- 1. JRA-55 Reanalysis system
 - Data assimilation system and boundary
 - Observation
 - Stream and progress
- 2. Early result





1. JRA-55 Reanalysis System



JRA-55 Reanalysis system



	JRA-25	JRA-55
Reanalysis years	1979-2004 (26 years)	1958-2012 (55 years)
Equivalent operational NWP system	As of Mar. 2004	As of Dec. 2009
Resolution	T106L40 (~120km) (top layer at 0.4 hPa)	TL319L60 (~60km) (top layer at 0.1 hPa)
Time integration	Eularian	Semi-Lagrangian
Assimilation scheme	3D-Var	4D-Var (with T106 inner model)
Bias correction (satellite radiance)	Adaptive method (Sakamoto et al. 2009)	Variational Bias Correction (Dee et al. 2009)
Tropical Cyclone	Wind profile retrievals (TCRs) provided by Dr.Fiorino were assimilated.	Same as JRA-25




Boundary and forcing fields



	JRA-25	JRA-55
Radiatively active gases	H ₂ O, CO ₂ , O ₃	H ₂ O, CO ₂ , O ₃ , CH ₄ , N ₂ O, CFC-11, CFC-12, HCFC-22
GHG concentrations	Constant at 375 ppmv (CO ₂)	Annual mean data are interpolated to daily data (CO2,CH4,N2O)
Ozone	Daily 3-D ozone (produced by AED/JMA)	(-1978) Monthly climatology (1979-) New daily 3-D ozone (produced using a revised CTM)
Aerosols	Annual climatology for continental and maritime aerosols	Monthly climatology for continental and maritime aerosols
SST Sea ice	COBE SST (Ishii <i>et al.</i> , 2005, <i>I.J.Clim.</i>)	COBE SST (ver. 1.5)




Observational data used in JRA-55





Available Reprocessed AMV and CSR data




Thick line: reprocessed period



JRA-55 progress status







Completed as of 2 April, 2012

JRA-55 will be completed in the first half of 2013.






2. Early result

Red line is JRA-55 in the following graphs. Note that only completed years are plotted.



Improvement of vertical temperature profiles






Vertical profiles of global mean bias and RMS difference between radiosonde temperature measurements and the background (solid lines) / analyzed fields (dotted lines) from JRA-25 (black) and JRA-55 (red) in January 1981.



Surface (2m) temperature



1982

2-m temperature anomalies, NH

Reanalysis - CRUTEM Ver. 3

NRA1 - CRU — JRA-25 - CRU ERA-40 - CRU — JRA-55 - CRU

Year


2000

1994

2006

2012

2-m temperature anomalies, SH




-0.4 L-1958

1964

1970

1976









Land Surface (2m) Relative Humidity

2-m relative humidity anomalies, NHland



JRA - HadCRUH

JRA-55 is better than JRA-25.







JRA-25 - HadCRUH — JRA-55 - HadCRUH



Zonal Mean Precipitation










Water budget in Amazon





←Good agreement with GPCP


In JRA-25, *Unrealistic dry bias* is found over the Amazon River basin.




RMSE(m) Z500 Northern Hemisphere ft=48

Comparison of Forecast Scores

RMSE of Z500 (48-hour forecast) for NH and SH









JRA-55 reference



Ebita et al. 2011

- Ayataka Ebita, Shinya Kobayashi, Yukinari Ota, Masami Moriya,
 Ryoji Kumabe, Kazutoshi Onogi, Yayoi Harada, Soichiro Yasui,
 Kengo Miyaoka, Kiyotoshi Takahashi, Hirotaka Kamahori, Chiaki
 Kobayashi, Hirokazu Endo, Motomu Soma, Yoshinori Oikawa and
 Takahisa Ishimizu
- "The Japanese 55-year Reanalysis "JRA-55": An Interim Report", SOLA, Vol. 7, pp.149-152 (2011) .
- http://www.jstage.jst.go.jp/article/sola/7/0/7_149/_article

JRA-25 reference (Onogi et al. 2007, JMSJ)

- http://www.jstage.jst.go.jp/article/jmsj/85/3/85 369/ article
- So far, about 2,000 registered users from 66 countries





Thank you for your attention

